

<https://helda.helsinki.fi>

---

## Developmental expression patterns of six6: a gene linked with spawning ecotypes in Atlantic salmon

Moustakas-Verho, Jacqueline

2020-09-30

---

Moustakas-Verho , J , Kurko , J , House , A , Erkinaro , J , Debes , P & Primmer , C 2020 , ' Developmental expression patterns of six6: a gene linked with spawning ecotypes in Atlantic salmon ' , Gene expression patterns , vol. 38 . <https://doi.org/10.1016/j.gep.2020.119149>

---

<http://hdl.handle.net/10138/319863>

<https://doi.org/10.1016/j.gep.2020.119149>

---

cc\_by\_nc\_nd

publishedVersion


---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*



# Developmental expression patterns of *six6*: A gene linked with spawning ecotypes in Atlantic salmon

Jacqueline Emmanuel Moustakas-Verho<sup>a,b,\*</sup>, Johanna Kurko<sup>a,b</sup>, Andrew H. House<sup>a,b</sup>,  
Jaakko Erkinaro<sup>c</sup>, Paul Debes<sup>a,b,1</sup>, Craig Robert Primmer<sup>a,b,\*\*</sup>

<sup>a</sup> Organismal and Evolutionary Biology Research Programme, University of Helsinki, Viikinkaari 9, 00014, Helsinki, Finland

<sup>b</sup> Institute of Biotechnology, University of Helsinki, Finland

<sup>c</sup> Natural Resources Institute Finland (LUKE), Oulu, Finland

## ARTICLE INFO

### Keywords:

Sine oculis homeobox  
*Six6*  
Atlantic salmon  
Development  
Sensory organs  
In situ  
Hybridization

## ABSTRACT

The Atlantic salmon has been studied extensively, particularly as a model for understanding the genetic and environmental contributions to the evolution and development of life history traits. Expression pattern analysis *in situ*, however, is mostly lacking in salmon. We examine the embryonic developmental expression of *six6*, a candidate gene previously identified to be associated with spawning ecotypes and age at sexual maturity, in Atlantic salmon. *Six6* is a member of the sine oculis homeobox family of transcription factors and is known to regulate eye and brain development in other vertebrates. We assay the expression of this gene in embryonic Atlantic salmon *Salmo salar* by whole-mount *in situ* hybridization. In line with earlier studies in other vertebrate species, we find conserved expression in the developing brain and sensory organs, including optic and olfactory primordia. However, we also find previously unreported domains of expression that suggest additional roles in axial and appendicular development, cardiovascular, intestinal, and sensory organogenesis. Each of these systems are important in the sensory ecology of Atlantic salmon, suggesting it is plausible that *six6* may have pleiotropic roles in this complex phenotype.

## 1. Introduction

The Atlantic salmon (*Salmo salar* L.) is an exciting system for exploring the genetic variation that underlies the evolution of phenotypic diversity. Atlantic salmon have an anadromous life cycle whereby mature fish spawn in rivers and juveniles remain in the freshwater environment for 1–7 years, after which they migrate to sea for a varying number of years before returning to reproduce, often in their natal river. Some of the traits behind this complex life-history strategy include adaptation to the local environments of their spawning grounds, variation in age at maturity, and the ability to transform their physiology for marine and freshwater environments (reviewed in Jonsson and Jonsson, 2011).

Recent studies examining patterns of variation across the genome of salmonids identified several genome regions inferred to be under diversifying selection and/or linked with the abovementioned traits (Bourret et al., 2013; Moore et al., 2014; Barson et al., 2015; Lemay and

Russello, 2015; Veale and Russello, 2017; Pritchard et al., 2018). One such locus, near the sine oculis-related homeobox 6 (*six6*) gene on chromosome 9, has been found to associate with age at maturity (Johnston et al., 2014; Barson et al., 2015; Sinclair-Waters et al., 2020) and spawning ecotype in salmonids, including Atlantic salmon and sockeye salmon, *Oncorhynchus nerka*. More specifically, *six6* alleles have been associated with stream vs lake-shore spawning ecotypes in sockeye salmon (Veale and Russello, 2017) and seasonal return migration timing of Atlantic salmon sub-populations (Pritchard et al., 2018; Cauwelier et al., 2018). *SIX6* also associates with size and age at maturity in humans (Perry et al., 2014; Day et al., 2017) and is required for fertility and puberty in mice (Larder et al., 2011; Pandolfi et al., 2019). In Atlantic salmon, the chromosome 9 *six6* paralogue is expressed in the brain, eye, gill, and testis, and is hypothesized to contribute to maturation regulation via the brain-pituitary-gonadal axis (Kurko et al., 2020).

*Six6* is a member of the sine oculis homeobox family of transcription

\* Corresponding author. Organismal and Evolutionary Biology Research Programme, University of Helsinki, Viikinkaari 9, 00014, Helsinki, Finland.

\*\* Corresponding author. Organismal and Evolutionary Biology Research Programme, University of Helsinki, Viikinkaari 9, 00014, Helsinki, Finland.

E-mail addresses: [jacqueline.moustakas@helsinki.fi](mailto:jacqueline.moustakas@helsinki.fi) (J.E. Moustakas-Verho), [craig.primmer@helsinki.fi](mailto:craig.primmer@helsinki.fi) (C.R. Primmer).

<sup>1</sup> Present address: Department of Aquaculture and Fish Biology, Hólar University College, Háeyri 1, 551 Sauðárkrúkur, Iceland.

factors that was first identified from a mutation in a gene called *sine oculis* in *Drosophila melanogaster* that affects the visual system (Cheyette et al., 1994; Serikaku and O'Tousa, 1994). *Six6* has evolutionary conserved roles in the control of eye development in fly, fish, frog, chick, and mouse (Seo et al., 1998; Toy et al., 1998; Jean et al., 1999; Lopez-Rios et al., 1999, 2003; Toy and Sundin, 1999; Zuber et al., 1999; Ghanbari et al., 2001; Ogawa et al., 2019). Two additional SIX family member genes are found in flies, *optix* and *DSix4*, and there are currently three subclasses of SIX proteins, each of which contain one fly gene and their orthologs: *Six1/2* (*sine oculis*), *Six3/6* (*optix*), and *Six4/5* (*DSix4*). *Optix* and its orthologs also regulate eye development, whereas *DSix4* and its orthologs function primarily in mesoderm derivatives (reviewed in Kumar 2009). SIX genes are broadly conserved in sequence and function, and frequently cooperate with other SIX genes to regulate organogenesis (Diacou et al., 2018; Garcez et al., 2014; Kawakami et al., 2000; Kobayashi et al., 2007; Ogawa et al., 2019). Despite the wealth of knowledge of expression patterns in the gene family, there is no

information about specific expression domains in salmon. Given the gene's link with ecologically relevant phenotypes, this would be an important first step towards understanding the possible functional mechanisms resulting in such ecologically relevant variation.

Here, we examine the developmental expression of *six6* in Atlantic salmon embryos from approximately the eyed-egg stage of development through hatching using whole-mount *in situ* hybridization. Our data support a conserved role for this gene in the development of the cranial sensory organs, and suggest the evolution of novel expression in Atlantic salmon potentially related to spawning and migration.

## 2. Results


The common ancestor of salmonids underwent a salmonid-specific fourth vertebrate whole-genome duplication (WGD) (Macqueen and Johnston, 2014; Lien et al., 2016), and, consequently, there are four paralogues of *six6* in the Atlantic salmon genome on chromosomes 1, 7,



**Fig. 1.** (A) Maximum-likelihood tree of SIX3/6 nucleotide sequences. Accession numbers are bracketed after each paralogue. (B) Amino acid alignment of SIX3 and SIX6 paralogues. The SIX domain (top) and homeodomain (bottom) sequences of SIX proteins are highly conserved.

C! (B& C ( C<K &(IC C?B &(IC C?B &(IC C?G 3 \* &(IC C!C  
1,J 7 1 8 7 % 8 &8 & ;<<?G " & ;<<?) & ;<< 8 &  
( <<K : (B 7 7 &8 & 7 8!  
" ( " C 7 7 &8!  
5 G & 3 " 6 ! (C C?B &(C C?G 3 \* &(C C?B &!  
48 "8 2 ; ! " !<  
70 & ;< ( 8 & " !< : 8 &  
, " ! (AB 7 • ! (C B?@ & ;< ( 8 & !  
" ! " ! " ! " ! 7 ! (C : 8 &  
7G (C C?B & ! " ! " ! " ! 9 ! 7% & !  
(C C?B & ! " ! " ! 6 & ;< < 8 & !  
7 & ) 8 & ! 7 & ) 8 & ! " !<  
(:B ! ! " 7 & / % 8 & (C @! !<  
" ! " " ! " " ! " ! " !<  
7 & - . 8 & ;:: ;AC 3 ! ! 7 ! ;< ( 8 &  
7 & L 1 ! ; ! ; J 8 & 9 ! ! ! 6 ! 6 ! 3 " 6 0 2 ! "  
;:: 7 & \* 8 & " ! 6 & 3 " 6 ! 75 " !<  
7 & 5 8 & 3 " 6 ! & ;< < ? G " & ;< < ? + + ;< < 8 & "  
48 4 8 2 2 2 5 & (AB ! " & 71 ;< < 8 !  
6 ! ! ! " ! " ! " ! " ! " ! " !<  
7 8 ! " ! 7 & ! ; 8 & ! " ! " ! " ! " ! " !<  
;:: 7 & / ! ; \$ ! - ! ; . L ! ; 1 D 8 & " ! " ! " ! " ! " ! " !<  
;AC ! ! ! 6 7 & ;< < ? & ;< ( ? 1 &(IC C?B &  
7 & 8 & (:B ! ! ! 6 7 + " ! " ! " ! " ! " ! " !<  
; , 8 & (C @ ;:: ! 6 7 L ! (C C?B 1 " !<  
7 & - ; 0 \* ! ; + D 8 & ; AC ! ! ;< ( 8 & " 6 ! " !<  
6 ! ! 6 7% 5 " ! (CA? L !<  
7 & ; M 1 8 & 6 (C C?B & 3 71 !<  
7 & M 8 & 3 + " (& ! ;< < 8 !<  
5 " " " 3 , & ! ;< < 8 & \* % 74  
" ! ! " ! " ! " ! " ! " ! " ! " ! " ! " ! " !<  
6 ! ! ! ! ! 6 6 " ! - " " 7% ' G 8  
! ! ! ! ! 6 6 " 7- & ;< ( ? ' 6 & ;< ( 8 & G \* %<  
& " !<  
3 " 6 ! ! 0 2 ! ! " !<  
! 71

- 9 -20.9339 Tm [( Tf 1 ) ( 1 47.1316 -11.7437 Tm [( ) ] TJ 1 0 0 1 66.9



**Fig. 2.** Developmental expression of *six3* in Atlantic salmon embryos. (A–E) 158  $\tau_s$  embryos shown in lateral (A, D), frontal (B), and dorsal (C) view, and in section (E). (F–K) 178  $\tau_s$  embryos shown in lateral (F, J), frontal (G, I), and dorsal (H) view, and in section (K). (L–P) 194  $\tau_s$  embryos shown in lateral (L, O), frontal (M), and dorsal (N) view, and in section (P). (Q–U) 227  $\tau_s$  embryos shown in lateral (Q, T), frontal (R), and dorsal (S) view, and in section (U). (V–Z) 259  $\tau_s$  embryos shown in lateral (V), dorsal (W–X), and ventral (Y) view, and in section (Z). af, anal fin; ax, axial vasculature; c, cerebellum; cf, caudal fin; cv, caudal vein; da, dorsal artery; ff, fin fold; g, gut; gi, gill; h, heart; he, hematopoietic; j, jaw; l, lateral line; m, myotome; ms, myosepta; n, notochord; nt, neural tube; ol, olfactory (black arrows); op, optic; ot, otic; p, pineal gland (white arrowhead); pf, pectoral fin; pv, periventricular tissue; r, retina; sf, slow muscle fibers; t, tectum (black arrowhead).

/  
 /  
 " ,&  
 ! 6 6  
 &  
 !  
 !  
 7) &!< ( ? \* !< ( :? ' &!<  
 ;< ( B & &7 ;< ( B 8 3  
 " "  
 = !  
 &  
 " 80  
 " "  
 ! ! !  
 &5  
 " ! ! !  
 ! ! !  
 ! ! !  
 0 12 3 42  
 1 6  
 1 , :: B ;< ( B &  
 7 8  
 . \* , /  
 7 2 ; @A 00 ! < ; AA : ( A 0 ! - !/ 80&  
 7 B & :) 8 ! ! !  
 &  
 7 8  
 7G (!C C82&  
 " & "  
 7 ' 4 1 86 @ K  
 (<<K ;< ) &  
 3 +  
 C ! G 5 1 74 &!< ( ? &!< ( ? 1  
 \* ;!< ( :? ) &!< ;< 8 ! ;< " 6 3  
 5 7 - . 8 " 7 85& " 7 8  
 / / " / 7 / 8 - . " 7 8  
 / " / ! " 7 \* 8 1  
 &!C C!A 7 ;< ( ( 8 6 " 3 2 < ) ! " !< ( &+< K'  
 A 4 ' 7 \* 8 ! 6 - 1 J - 1 C &  
 ' 4 1 - 1 J 1 , J " G 4 7 == &  
 1 " & & & = 8 " &G = ' 7 ;< ;< & ( ! == &  
 " & 8 8 1 , J &!< ( 8 & - 7  
 ) 0 % \$ M B A " 7  
 - &!< < 8  
 1 , J 7\$ ;< < 8 &  
 !  
 !  
 " &  
 " "  
 " "  
 7 &!< ;< 8 &  
 3 !  
 7 &!< ;< 8 &



## Author contributions

Jacqueline Moustakas-Verho: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Visualization, Writing - original draft, Writing - review & editing. Johanna Kurko: Conceptualization, Data curation, Investigation, Methodology, Writing - review & editing. Andrew House: Data curation, Investigation, Writing - review & editing. Jaakko Erkinaro: Resources. Paul Debes: Conceptualization, Data curation, Investigation, Methodology. Craig Primmer: Conceptualization, Data curation, Funding acquisition, Investigation, Resources, Writing - review & editing. All authors approved the manuscript.

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

We thank Suvi Ikonen, Outi Ovaskainen, Spiros Papakostas, Victoria Pritchard, Annukka Ruokolainen, Marion Sinclair-Waters, Ksenia Zueva, and LUKE Laukaa staff for help related to gamete stripping and fish husbandry, Outi Kostia for technical advice, and Jukka-Pekka Verta for advice with Geneious.

## References

- Aykanat, T., Rasmussen, M., Ozerov, M., Niemelä, E., Paulin, L., Vähä, J.-P., Hindar, K., Wennevik, V., Pedersen, T., Svenning, M.-A., Primmer, C., 2020. Life-history genomic regions explain differences in Atlantic salmon marine diet specialization. *J. Animal Ecol.* <https://doi.org/10.1111/1365-2656.13324>.
- Barson, N.J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G.H., Fiske, P., Jacq, C., Jensen, A.J., Johnston, S.E., Karlsson, S., Kent, M., Moen, T., Niemelä, E., Nome, T., Naesje, T.F., Orell, P., Romakkaniemi, A., Saegrov, H., Urdal, K., Erkinaro, J., Lien, S., Primmer, C.R., 2015. Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature* 528 (7582), 405–408.
- Bourret, V., Kent, M.P., Primmer, C.R., Vasemägi, A., Karlsson, S., Hindar, K., McGinnity, P., Verspoor, E., Bernatchez, L., Lien, S., 2013. SNP-array reveals genome-wide patterns of geographical and potential adaptive divergence across the natural range of Atlantic salmon (*Salmo salar*). *Mol. Ecol.* 22 (3), 532–551.
- Broglio, C., Rodríguez, F., Salas, C., 2003. Spatial cognition and its neural basis in teleost fishes. *Fish Fish.* 4 (3), 247–255.
- Brugmann, S.A., Moody, S.A., 2005. Induction and specification of the vertebrate ectodermal placodes: precursors of the cranial sensory organs. *Biol. Cell.* 97 (5), 303–319.
- Cauwelier, E., Gilbey, J., Sampayo, J., Stradmeyer, L., Middlemas, S.J., 2018. Identification of a single genomic region associated with seasonal river return timing in adult Scottish Atlantic salmon (*Salmo salar*), using a genome-wide association study. *Can. J. Fish. Aquat. Sci.* 75 (9), 1427–1435.
- Cheyette, B.N., Green, P.J., Martin, K., Garren, H., Hartenstein, V., Zipursky, S.L., 1994. The *Drosophila sine oculis* locus encodes a homeodomain-containing protein required for the development of the entire visual system. *Neuron* 12 (5), 977–996.
- Conte, I., Morcillo, J., Bovolenta, P., 2005. Comparative analysis of *Six3* and *Six6* distribution in the developing and adult mouse brain. *Dev. Dynam.* 234 (3), 718–725.
- Day, F.R., Thompson, D.J., Helgason, H., Chasman, D.I., Finucane, H., Sulem, P., et al., 2017. Genomic analyses identify hundreds of variants associated with age at menarche and support a role for puberty timing in cancer risk. *Nat. Genet.* 49 (6), 834–841.
- Diacou, R., Zhao, Y., Zheng, D., Cvekl, A., Liu, W., 2018. *Six3* and *six6* are jointly required for the maintenance of multipotent retinal progenitors through both positive and negative regulation. *Cell Rep.* 25 (9), 2510–2523.
- Dittman, A., Quinn, T., 1996. Homing in Pacific salmon: mechanisms and ecological basis. *J. Exp. Biol.* 199 (1), 83–91.
- Ekström, P., van Veen, T., 1984. Pineal neural connections with the brain in two teleosts, the crucian carp and the European eel. *J. Pineal Res.* 1 (3), 245–261.
- Fenwick, J.C., 1970. The pineal organ. In: *Fish Physiology*, vol. 4. Academic Press, pp. 91–108.
- Garcez, R.C., Douarin, Le, M. N., Creuzet, S.E., 2014. Combinatorial activity of *Six1-2-4* genes in cephalic neural crest cells controls craniofacial and brain development. *Cell. Mol. Life Sci.* : CMLS 71 (11), 2149–2164.
- Ghanbari, H., Seo, H.C., Fjose, A., Brändli, A.W., 2001. Molecular cloning and embryonic expression of *Xenopus Six* homeobox genes. *Mech. Dev.* 101 (1–2), 271–277.
- Gorodilov, Y.N., 1996. Description of the early ontogeny of the Atlantic salmon, *Salmo salar*, with a novel system of interval (state) identification. *Environ. Biol. Fish.* 47 (2), 109–127.
- Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* 59 (3), 307–321.
- Hasegawa, M., Kishino, H., Yano, T., 1985. Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. *J. Mol. Evol.* 22, 160–174.
- Hasler, A.D., Wisby, W.J., 1951. Discrimination of stream odors by fishes and its relation to parent stream behavior. *Am. Nat.* 85 (823), 223–238.
- Holmqvist, B.I., Östholm, T., Ekström, P., 1994. Neuroanatomical analysis of the visual and hypophysiotropic systems in Atlantic salmon (*Salmo salar*) with emphasis on possible mediators of photoperiodic cues during parr-smolt transformation. *Aquaculture* 121 (1–3), 1–12.
- Jayne, B.C., Lauder, G.V., 1994. How swimming fish use slow and fast muscle fibers: implications for models of vertebrate muscle recruitment. *J. Comp. Physiol.* 175 (1), 123–131.
- Jean, D., Bernier, G., Gruss, P., 1999. *Six6* (*Optx2*) is a novel murine *Six3*-related homeobox gene that demarcates the presumptive pituitary/hypothalamic axis and the ventral optic stalk. *Mech. Dev.* 84 (1–2), 31–40.
- Johnston, S.E., Orell, P., Pritchard, V.L., Kent, M.P., Lien, S., Niemelä, E., Erkinaro, J., Primmer, C.R., 2014. Genome-wide SNP analysis reveals a genetic basis for sea-age variation in a wild population of Atlantic salmon (*Salmo salar*). *Mol. Ecol.* 23, 3452–3468.
- Jonsson, Bror, Jonsson, Nina, 2011. Ecology of Atlantic salmon and Brown Trout: Habitat as A Template For Life Histories. Springer, Dordrecht.
- Kawakami, K., Sato, S., Ozaki, H., Ikeda, K., 2000. Six family genes—structure and function as transcription factors and their roles in development. *Bioessays* 22 (7), 616–626.
- Kobayashi, M., Osanai, H., Kawakami, K., Yamamoto, M., 2000. Expression of three zebrafish *Six4* genes in the cranial sensory placodes and the developing somites. *Mech. Dev.* 98 (1–2), 151–155.
- Kobayashi, H., Kawakami, K., Asashima, M., Nishinakamura, R., 2007. *Six1* and *Six4* are essential for *Gdnf* expression in the metanephric mesenchyme and ureteric bud formation, while *Six1* deficiency alone causes mesonephric-tubule defects. *Mech. Dev.* 124 (4), 290–303.
- Kryvi, H., Rusten, I., Fjellidal, P.G., Nordvik, K., Totland, G.K., Karlsen, T., et al., 2017. The notochord in Atlantic salmon (*Salmo salar* L.) undergoes profound morphological and mechanical changes during development. *J. Anat.* 231 (5), 639–654.
- Kumar, J.P., 2009. The sine oculis homeobox (*SIX*) family of transcription factors as regulators of development and disease. *Cell. Mol. Life Sci.* 66 (4), 565.
- Kurko, J., Debes, P.V., House, A.H., Aykanat, T., Erkinaro, J., Primmer, C.R., 2020. Transcription profiles of age-at-maturity-associated genes suggest cell fate commitment regulation as a key factor in the Atlantic salmon maturation process. *G3: Genes, Genomes, Genetics* 10 (1), 235–246.
- Larder, R., Clark, D.D., Miller, N.L., Mellon, P.L., 2011. Hypothalamic dysregulation and infertility in mice lacking the homeodomain protein. *Six6*. *J. Neurosci.* 31 (2), 426–438.
- Lemay, M.A., Russello, M.A., 2015. Genetic evidence for ecological divergence in kokanee salmon. *Mol. Ecol.* 24 (4), 798–811.
- Li, X., Perissi, V., Liu, F., Rose, D.W., Rosenfeld, M.G., 2002. Tissue-specific regulation of retinal and pituitary precursor cell proliferation. *Science* 297 (5584), 1180–1183.
- Lien, S., Koop, B.F., Sandve, S.R., Miller, J.R., Kent, M.P., Nome, T., et al., 2016. The Atlantic salmon genome provides insights into rediploidization. *Nature* 533 (7602), 200–205.
- López-Ríos, J., Gallardo, M.E., de Córdoba, S.R., Bovolenta, P., 1999. *Six9* (*Optx2*), a new member of the six gene family of transcription factors, is expressed at early stages of vertebrate ocular and pituitary development. *Mech. Dev.* 83 (1–2), 155–159.
- López-Ríos, J., Tessmar, K., Loosli, F., Wittbrodt, J., Bovolenta, P., 2003. *Six3* and *Six6* activity is modulated by members of the groucho family. *Development* 130 (1), 185–195.
- Macqueen, D.J., Robb, D., Johnston, I.A., 2007. Temperature influences the coordinated expression of myogenic regulatory factors during embryonic myogenesis in Atlantic salmon (*Salmo salar* L.). *J. Exp. Biol.* 210 (16), 2781–2794.
- Macqueen, D.J., Johnston, I.A., 2014. A well-constrained estimate for the timing of the salmonid whole genome duplication reveals major decoupling from species diversification. *Proc. Biol. Sci.* 281 (1778), 20132881.
- Maddison, W.P., Maddison, D.R., 2019. Mesquite: a Modular System for Evolutionary Analysis, Version 3.61. <http://www.mesquiteproject.org>.
- Mano, H., Asaoka, Y., Kojima, D., et al., 2019. Brain-specific homeobox *Bsx* specifies identity of pineal gland between serially homologous photoreceptive organs in zebrafish. *Commun. Biol.* 2, 364. <https://doi.org/10.1038/s42003-019-0613-1>.
- Marchesan, M., Ota, D., Ferrero, E.A., 2000. The role of mechanical stimulation during breeding in the grass goby *Zosterisessor ophiocephalus* (Teleostei, Gobiidae). *Ital. J. Zool.* 67 (1), 25–30.
- Medina, L.M., Garcia, C.M., Urbina, A.F., Manjarrez, J., Moyaho, A., 2013. Female vibration discourages male courtship behaviour in the Amarillo fish (*Girardinichthys multiradiatus*). *Behav. Process.* 100, 163–168.
- Moore, J.S., Bourret, V., Dionne, M., Bradbury, I., O'Reilly, P., Kent, M., et al., 2014. Conservation genomics of anadromous Atlantic salmon across its North American range: outlier loci identify the same patterns of population structure as neutral loci. *Mol. Ecol.* 23 (23), 5680–5697.
- Motta, M., Fraschini, F., Martini, L., 1967. Endocrine effects of pineal gland and of melatonin. *PSEBM (Proc. Soc. Exp. Biol. Med.)* 126 (2), 431–435.

- Moustakas-Verho, J.E., McLennan, R., Spengler, J., Kulesa, P.M., Cebra-Thomas, J.A., 2019. Manipulation of developmental function in turtles with notes on alligators. In: *Vertebrate Embryogenesis*. Humana Press, New York, NY, pp. 247–263.
- Northmore, D., 2011. Optic tectum. *Encyclop. Fish Physiol.: From Genome Environ.* 1, 131–142.
- Ogawa, Y., Shiraki, T., Asano, Y., Muto, A., Kawakami, K., Suzuki, Y., et al., 2019. *Six6* and *Six7* coordinately regulate expression of middle-wavelength opsins in zebrafish. *Proc. Natl. Acad. Sci. Unit. States Am.* 116 (10), 4651–4660.
- Olivares, J., Schmachtenberg, O., 2019. An update on anatomy and function of the teleost olfactory system. *PeerJ* 7, e7808.
- Pandolfi, E.C., Tonsfeldt, K.J., Hoffmann, H.M., Mellon, P.L., 2019. Deletion of the homeodomain protein *Six6* from GnRH neurons decreases GnRH gene expression, resulting in infertility. *Endocrinology* 160 (9), 2151–2164.
- Perry, J.R., Day, F., Elks, C.E., Sulem, P., Thompson, D.J., Ferreira, T., et al., 2014. Parent-of-origin-specific allelic associations among 106 genomic loci for age at menarche. *Nature* 514 (7520), 92–97.
- Pritchard, V.L., Mäkinen, H., Vähä, J.P., Erkinaro, J., Orell, P., Primmer, C.R., 2018. Genomic signatures of fine-scale local selection in Atlantic salmon suggest involvement of sexual maturation, energy homeostasis and immune defence-related genes. *Mol. Ecol.* 27 (11), 2560–2575.
- Satou, M., Takeuchi, H.A., Nishii, J., Tanabe, M., Kitamura, S., Okumoto, N., Iwata, M., 1994. Behavioral and electrophysiological evidences that the lateral line is involved in the inter-sexual vibrational communication of the himé salmon (landlocked red salmon, *Oncorhynchus nerka*). *J. Comp. Physiol.* 174 (5), 539–549.
- Schlosser, G., 2006. Induction and specification of cranial placodes. *Dev. Biol.* 294 (2), 303–351.
- Seo, H.C., Drivenes, Ø., Ellingsen, S., Fjose, A., 1998. Transient expression of a novel *Six3*-related zebrafish gene during gastrulation and eye formation. *Gene* 216 (1), 39–46.
- Serikaku, M.A., O'Tousa, J.E., 1994. *Sine oculis* is a homeobox gene required for *Drosophila* visual system development. *Genetics* 138 (4), 1137–1150.
- Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., Söding, J., Thompson, J.D., Higgins, D.G., 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol. Syst. Biol.* 7, 539.
- Sinclair-Waters, M., Ødegård, J., Korsvoll, S.A., Moen, T., Lien, S., Primmer, C.R., Barson, N.J., 2020. Beyond large-effect loci: large-scale GWAS reveals a mixed large-effect and polygenic architecture for age at maturity of Atlantic salmon. *Genet. Sel. Evol.* 52 (1), 1–11.
- Stemple, D.L., 2005. Structure and function of the notochord: an essential organ for chordate development. *Development* 132 (11), 2503–2512.
- Sweetman, D., 2011. In situ detection of microRNAs in animals. In: *MicroRNAs in Development*. Humana Press, pp. 1–8.
- Thisse, B., Thisse, C., 2004. Fast release clones: a high throughput expression analysis. ZFIN Direct Data Submission 2004.
- Toy, J., Yang, J.M., Leppert, G.S., Sundin, O.H., 1998. The *optx2* homeobox gene is expressed in early precursors of the eye and activates retina-specific genes. *Proc. Natl. Acad. Sci. Unit. States Am.* 95 (18), 10643–10648.
- Toy, J., Sundin, O.H., 1999. Expression of the *optx2* homeobox gene during mouse development. *Mech. Dev.* 83 (1–2), 183–186.
- Valente, L.M., Moutou, K.A., Conceição, L.E., Engrola, S., Fernandes, J.M., Johnston, I.A., 2013. What determines growth potential and juvenile quality of farmed fish species? *Rev. Aquacult.* 5, S168–S193.
- Veale, A.J., Russello, M.A., 2017. An ancient selective sweep linked to reproductive life history evolution in sockeye salmon. *Sci. Rep.* 7 (1), 1–10.
- Zhou, S., Ackman, R.G., Morrison, C., 1995. Storage of lipids in the myosepta of Atlantic salmon (*Salmo salar*). *Fish Physiol. Biochem.* 14 (2), 171–178.
- Zhou, S., Ackman, R.G., Morrison, C., 1996. Adipocytes and lipid distribution in the muscle tissue of Atlantic salmon (*Salmo salar*). *Can. J. Fish. Aquat. Sci.* 53 (2), 326–332.
- Zuber, M.E., Perron, M., Philpott, A., Bang, A., Harris, W.A., 1999. Giant eyes in *Xenopus laevis* by overexpression of *XOptx2*. *Cell* 98 (3), 341–352.